

On Neutrino Masses and a Low Breaking Scale of Left-Right Symmetry

Oleg Khasanov and Gilad Perez

Department of Particle Physics, Weizmann Institute of Science, Rehovot 76100, Israel

Abstract

In left-right symmetric models (LRS) the light neutrino masses arise from two sources: the seesaw mechanism and a VEV of an $SU(2)_L$ triplet. If the left-right symmetry breaking, v_R , is low, $v_R \lesssim 15$ TeV, the contributions to the light neutrino masses from both the seesaw mechanism and the triplet Yukawa couplings are expected to be well above the experimental bounds. We present a minimal LRS with an additional $U(1)$ symmetry in which the masses induced by the two sources are below the eV scale and the two-fold problem is solved. We further show that, if the $U(1)$ symmetry is also responsible for the lepton flavor structure, the model yields a small mixing angle within the first two lepton generations.

I. INTRODUCTION

Among the various new physics (NP) models that give neutrino masses, the left-right symmetric (LRS) framework is, in many senses, one of the most attractive. A left-right symmetric model (LRS) is based on the gauge group $SU(2)_R \times SU(2)_L \times U(1)_{B-L}$ [1]. At the high energy scale of the LRS breaking, v_R , various new fields and interactions are present. In many LRS, *e.g.* LRS embedded in GUT models [2], the scale v_R is much higher than the electroweak (EW) breaking scale, k , and the low energy effective Lagrangian is similar in many aspects to that of the Standard Model (SM). In such cases present and near future experiments will not be able to directly probe the NP. We therefore investigate whether a natural LRS with $v_R \lesssim 15$ TeV can be constructed consistent with the experimental data on the lepton flavor parameters. In particular we require the active neutrino masses to be below the eV scale,

$$m_{\nu_i} \lesssim 1 \text{ eV} , \quad (1)$$

with $i = 1, 2, 3$. This is motivated by the cosmological [3], and direct [4,5] upper bounds on the active neutrino masses and in particular by the ones deduced from the atmospheric and solar experiments assuming hierarchical neutrino masses [6–8]. The charged lepton masses are [4],

$$m_e \simeq 0.51 \text{ MeV} , \quad m_\mu \simeq 105.7 \text{ MeV} , \quad m_\tau \simeq 1777 \text{ MeV} . \quad (2)$$

In section II we discuss a two-fold problem related to neutrino masses which one faces when dealing with a relatively low scale of v_R ; In section III we present a minimal LRSM (MLRSM) with an additional U(1) symmetry, which solves the above problem. In section IV we show that if the same U(1) symmetry accounts also for the lepton flavor structure such a model cannot yield a large mixing angle (LMA) between the first two generations. Comments and conclusions are given in section V.

II. NATURALNESS OF THE MSLRM WITH A LOW v_R

Below we investigate the consequences of having a low value for the LR symmetry breaking scale,

$$v_R \lesssim 15 \text{ TeV} . \quad (3)$$

The Lagrangian of the lepton sector in the MLRSM is given by:

$$\mathcal{L} = \mathcal{L}_0 + f \bar{L}_L \phi_1 L_R + g \bar{L}_L \phi_2 L_R + h(\Delta_L L_L L_L + \Delta_R L_R L_R) + V(\phi, \Delta) + h.c. , \quad (4)$$

where $\mathcal{L}_0$ contains the kinetic and the gauge interaction terms for the various fields. The field L is a lepton doublet, the ϕ field is a Higgs bi-doublet, $\phi_1 = \begin{pmatrix} h_1^0 & h_1^+ \\ h_2^- & h_2^0 \end{pmatrix}$, $\phi_2 = \tau_2 \phi_1^* \tau_2$, Δ_L [Δ_R] is a triplet of the SU(2)_L [SU(2)_R] gauge group and the scalar potential, $V(\phi, \Delta)$, is given below.

In the general case the scalar fields, $\Delta_{L,R}$ and ϕ , develop complex VEVs,

$$\langle \Delta_{L,R} \rangle = \begin{pmatrix} 0 & 0 \\ v_{L,R} e^{i\alpha_{L,R}} & 0 \end{pmatrix} , \quad \langle \phi_1 \rangle = \begin{pmatrix} k_1 e^{i\alpha_1} & 0 \\ 0 & k_2 e^{i\alpha_2} \end{pmatrix} . \quad (5)$$

In order to have a phenomenologically viable model the VEVs must be hierarchical,

$$|v_L| \ll k \ll |v_R| , \quad k = \sqrt{|k_1|^2 + |k_2|^2} \sim 250 \text{ GeV} . \quad (6)$$

If the ratio $r \equiv |k_2/k_1|$ is of order unity then in general no splitting is obtained between the Dirac masses of the two components of the SU(2)_L doublets. This causes phenomenological

problems both in the lepton sector (see discussion below) and in the quark sector [9–11]. Thus, we assume below that r is a small number.

Given the VEV hierarchy of eq. (6) the light neutrino mass matrix, M_l^ν , is given by:

$$M_l^\nu \approx (M_\nu^D)^T (M_R^{Maj})^{-1} M_\nu^D + M_L^{Maj}, \quad (7)$$

with M_ν^D being the Dirac neutrino mass and M_R^{Maj} [M_L^{Maj}] being the Majorana mass matrix for the right [left] handed neutrinos. The RHS of eq. (7) contains two terms. The first, $M_{See}^\nu \equiv (M_\nu^D)^T (M_R^{Maj})^{-1} M_\nu^D$, is related to the seesaw mechanism [12]. It depends on the scale of the Dirac neutrino masses and on v_R . The second, M_L^{Maj} , depends on v_L . In order to estimate the magnitude of M_l^ν we consider them separately.

A. The natural value of M_{See}^ν and M_L^{Maj}

1. M_{See}^ν

In general, without further assumptions, the scale of the charged lepton and the neutrino Dirac masses is roughly the same. Therefore, neglecting mixing, the value of the tau neutrino mass or $(M_{See}^\nu)_{33}$ is simply given by the well known seesaw formula [12]:

$$(M_{See}^\nu)_{33} \sim (m_\tau)^2 / v_R = \mathcal{O}(0.1 \text{ MeV}). \quad (8)$$

Such a high value is well above the bound given in eq. (1).

2. M_L^{maj}

The scalar potential of the MLRSM is given by ¹,

$$V(\phi, \Delta) = V_\phi + V_\Delta + V_{\phi\Delta}, \quad (9)$$

where

$$V_\phi = -\mu_{ij}^2 \text{Tr}(\phi_i^\dagger \phi_j) + \lambda_{ijkl} \text{Tr}(\phi_i^\dagger \phi_j) \text{Tr}(\phi_k^\dagger \phi_l), \quad (10)$$

$$V_\Delta = -\mu_i^2 \text{Tr}(\Delta_i \Delta_i^\dagger) + \rho_{ij} \text{Tr}(\Delta_i \Delta_i) \text{Tr}(\Delta_j^\dagger \Delta_j^\dagger), \quad (11)$$

with $\Delta_{1,2} = \Delta_{L,R}$ and

¹There is a redundancy in different terms in the potential the way it is written in eqs. (10-12). It does not, however, affect our analysis.

$$\begin{aligned}
V_{\phi\Delta} = & \alpha_{ij} \text{Tr}(\phi_i^\dagger \phi_j) \text{Tr}(\Delta_k \Delta_k^\dagger) + \text{Tr}(\beta_{ij} \Delta_L^\dagger \Delta_L \phi_i \phi_j^\dagger + \beta'_{ij} \Delta_R^\dagger \Delta_R \phi_i^\dagger \phi_j) \\
& + \gamma_{ij} \text{Tr}(\Delta_L^\dagger \phi_i \Delta_R \phi_j^\dagger).
\end{aligned} \tag{12}$$

It leads to the following relation [1,13–15]:

$$|v_R v_L| = a |k_1|^2 (1 + O(|r|)), \tag{13}$$

where a is a function of the various couplings. Without fine tuning a is expected to be of order unity. Substituting the values of the relevant VEVs given in eq. (3) and eq. (6) into eq. (13) leads to a rather high value of v_L ,

$$v_L = \mathcal{O}(1 \text{ GeV}). \tag{14}$$

Such a high value of v_L , with the triplet Yukawa couplings of order unity, yields light neutrino masses of the order of 1 GeV which is above the direct experimental bound [4,5].

To conclude, with $v_R \lesssim 15 \text{ TeV}$ and the Yukawas of the Higgs triplet of $\mathcal{O}(1)$, we expect $m_{\nu_\tau} = \mathcal{O}(0.1 - 10^3 \text{ MeV})$, which is well above the eV scale.

III. A MLRSM WITH A U(1) SYMMETRY MODEL

Below we show how an additional $U(1)$ symmetry can solve the problem of too massive active neutrinos discussed above. For our demonstration we consider a model with only third generation leptons. Since the seesaw mechanism relates the heaviest charged lepton to the heaviest neutrino, our model deals with the most severe phenomenological problem.

The low energy effective theory with the additional $U(1)$ symmetry is assumed to be broken by a small parameter, ε . Thus, various terms in the low energy effective Lagrangian are suppressed by powers of ε ,

$$\begin{aligned}
\mathcal{L}^{eff} = & \mathcal{L}_0 + \varepsilon^{|Q(L_R)-Q(L_L)+Q(\phi_1)|} f \bar{L}_L \phi_1 L_R + \varepsilon^{|Q(L_R)-Q(L_L)-Q(\phi_1)|} g \bar{L}_L \phi_2 L_R \\
& + \varepsilon^{|Q(\Delta_L)+2Q(L_L)|} h (\Delta_L L_L L_L + \Delta_R L_R L_R) + V(\phi, \Delta) + h.c.,
\end{aligned} \tag{15}$$

where we redefined the various Yukawa couplings so that $f, g, h = \mathcal{O}(1)$. The structure of $V(\phi, \Delta)$ can be easily deduced from eqs. (9-12). To our consideration it is important only to focus on the α_{ij}, β_{ij} and γ_{ij} terms. We redefine them so that now $\alpha_{ij}, \beta_{ij}, \gamma_{ij} = \mathcal{O}(1)$,

$$\alpha_{ij}, \beta'_{ji}, \beta_{ij} \Rightarrow \begin{cases} (\alpha_{ij}, \beta'_{ji}, \beta_{ij}) & i = j \\ (\alpha_{ij}, \beta'_{ji}, \beta_{ij}) \varepsilon^{2|Q(\phi_1)|} & i \neq j \end{cases} \tag{16}$$

and

$$\gamma_{ij} \Rightarrow \begin{cases} \gamma_{ij} \varepsilon^{|Q(\Delta_R)-Q(\Delta_L)|} & i = j \\ \gamma_{ij} \varepsilon^{|Q(\Delta_R)-Q(\Delta_L)+2Q(\phi_1)|} & i = 1 \quad j = 2 \\ \gamma_{ij} \varepsilon^{|Q(\Delta_R)-Q(\Delta_L)-2Q(\phi_1)|} & i = 2 \quad j = 1 \end{cases}. \tag{17}$$

The most general VEVs that the fields can develop are given in eq. (5). Using, however, $SU(2)_L \times SU(2)_R$ transformations of the form

$$U_{L,R} = \begin{pmatrix} e^{i\theta_{L,R}} & 0 \\ 0 & e^{-i\theta_{L,R}} \end{pmatrix} \quad (18)$$

one can bring v_L and v_R to be real. Thus, the effective potential acquires the following form at the minimum:

$$\begin{aligned} V(\langle\phi_{1,2}\rangle, \langle\Delta_{R,L}\rangle) = & -\mu_3^2(v_L^2 + v_R^2) + \frac{\rho}{4}(v_L^4 + v_R^4) + \frac{\rho'}{2}v_L^2v_R^2 + (v_L^2 + v_R^2)[(\alpha_{11} + \alpha_{22} + \bar{\beta}_{11})|k_1|^2 \\ & + (\alpha_{11} + \alpha_{22} + \bar{\beta}_{22})|k_2|^2 + \varepsilon^{|2Q(\phi_1)|}(4\alpha_{12} + 2\beta_{12})\mathcal{R}e(k_1k_2^*)] \\ & + 2v_Lv_R[\varepsilon^{|Q(\Delta_R)-Q(\Delta_L)|}(\gamma_{11} + \gamma_{22})\mathcal{R}e(k_1k_2^*) \\ & + \varepsilon^{|Q(\Delta_R)-Q(\Delta_L)+2Q(\phi_1)|}\gamma_{12}|k_1|^2 + \varepsilon^{|Q(\Delta_R)-Q(\Delta_L)-2Q(\phi_1)|}\gamma_{21}|k_2|^2] \\ & + \text{terms which depend only on } k_1, k_2, \end{aligned} \quad (19)$$

where $\rho = 4(\rho_{11} + \rho_{22})$, $\rho' = \rho_{12} + \rho_{21}$ and $\bar{\beta}_{ij} = \beta_{ij} + \beta'_{ij}$. As discussed above we assume $r, \varepsilon \ll 1$. Therefore the potential $V(\phi, \Delta)$ is approximately given by:

$$\begin{aligned} V(\langle\phi\rangle, \langle\Delta\rangle) \approx & -\mu_3^2(v_R^2 + v_L^2) + \frac{\rho}{4}(v_R^4 + v_L^4) + \frac{\rho'}{2}v_R^2v_L^2 \\ & + \frac{\alpha}{2}(v_R^2 + v_L^2)|k_1|^2 + \beta v_Lv_R|k_1|^2, \end{aligned} \quad (20)$$

with

$$\begin{aligned} \alpha \approx & 2(\alpha_{11} + \alpha_{22} + \bar{\beta}_{11}), \\ \beta \approx & 2[r^2\varepsilon^{|Q(\Delta_R)-Q(\Delta_L)+2Q(\phi_1)|}\gamma_{12} + \varepsilon^{|Q(\Delta_R)-Q(\Delta_L)-2Q(\phi_1)|}\gamma_{21} \\ & + r\cos(\alpha_1 - \alpha_2)\varepsilon^{|Q(\Delta_R)-Q(\Delta_L)|}(\gamma_{11} + \gamma_{22})]. \end{aligned} \quad (21)$$

Following the standard analysis of the minimization of the potential (see *e.g.* [1,13–16]) we obtain:

$$v_Lv_R = \gamma|k_1|^2, \quad (22)$$

with $\gamma = \beta/(\rho - \rho')$. Thus, in our model the suppression related to γ is roughly given by:

$$\gamma \sim \left\{ \max[r^2\varepsilon^{|Q(\Delta_L)-Q(\Delta_R)-2Q(\phi_1)|}, r\varepsilon^{|Q(\Delta_L)-Q(\Delta_R)|}, \varepsilon^{|Q(\Delta_L)-Q(\Delta_R)+2Q(\phi_1)|}] \right\}, \quad (23)$$

which can naturally bring v_L to below the eV scale as required by eq. (1).

In our model the lepton masses are given by:

$$\begin{aligned} m_{\nu_\tau}^D &= \frac{|k_1|}{\sqrt{2}}(f\varepsilon^{|Q(L_L)-Q(L_R)-Q(\phi_1)|} + rg\varepsilon^{|Q(L_L)-Q(L_R)+Q(\phi_1)|}e^{i\theta}) \\ m_\tau^D &= \frac{|k_1|}{\sqrt{2}}(g\varepsilon^{|Q(L_L)-Q(L_R)+Q(\phi_1)|} + rf\varepsilon^{|Q(L_L)-Q(L_R)-Q(\phi_1)|}e^{-i\theta}) \end{aligned} \quad (24)$$

where we used a phase convention in which v_R, k_1 are real.

To make our discussion concrete we choose the following set of charges:

$$Q(\Delta_L) = -Q(\Delta_R) = 6, \quad Q(\phi_1) = -Q(\phi_2) = 2, \quad Q(L_L) = -Q(L_R) = 3. \quad (25)$$

Assuming $\varepsilon^4 \sim r \sim 1/250$, $v_R \sim 15$ TeV, $k_1 \sim 250$ GeV we get

$$v_L \sim \varepsilon^{16} k_1^2 / v_R \sim 0.1 \text{ eV}, \quad (26)$$

$$m_\tau \sim \varepsilon^4 k_1 g \sim 1 \text{ GeV}, \quad m_{\nu_\tau}^D \sim \varepsilon^4 k_1 |\varepsilon^4 f + r g e^{i\theta}| \sim 1 \text{ MeV}, \quad (27)$$

and

$$m_{\nu_\tau} \sim h v_L - (m_{\nu_\tau}^D)^2 / v_R \sim 0.1 \text{ eV}. \quad (28)$$

To summarize, we showed that, in principle, using an additional U(1) symmetry and under the assumption that r is of $\mathcal{O}(10^{-2} - 10^{-3})$, one can construct a model in which v_R is of $\mathcal{O}(15 \text{ TeV})$, the neutrino masses are below $\mathcal{O}(1 \text{ eV})$ and $m_\tau \sim 1 \text{ GeV}$. In that sense, models in which the typical scale of LRS breaking is relatively low might still be natural.

IV. CAN A U(1) HORIZONTAL SYMMETRY ACCOUNT FOR THE RECENT SN DATA?

We examine below the possibility that the structure of the lepton flavor sector is explained solely by the above U(1) symmetry, which acts on the different flavors as an horizontal symmetry. In particular we focus on the mixing between the first two generations and investigate whether a large mixing angle (LMA) is obtained by the model, as strongly favored by the recent data from the SN experiments [6,8]. We assume for simplicity that all the right handed neutrinos are heavy. Thus our analysis does not apply to models of four light neutrinos. We also neglect CP violation (CPV) in the lepton sector.

The charged lepton mass matrix, M^{cl} is given by:

$$M^{cl} \sim \frac{k}{\sqrt{2}} \begin{pmatrix} \varepsilon^{|Q_3+Q_{13}^L+Q_{13}^R|} & \varepsilon^{|Q_3+Q_{23}^L+Q_{13}^R|} & \varepsilon^{|Q_3+Q_{13}^R|} \\ \varepsilon^{|Q_3+Q_{23}^R+Q_{13}^L|} & \varepsilon^{|Q_3+Q_{23}^L+Q_{23}^R|} & \varepsilon^{|Q_3+Q_{23}^R|} \\ \varepsilon^{|Q_3+Q_{13}^L|} & \varepsilon^{|Q_3+Q_{23}^L|} & \varepsilon^{|Q_3|} \end{pmatrix}, \quad (29)$$

with $Q_i = Q(L_R^i) - Q(L_L^i) + Q(\phi_1)$, $Q_{ij}^R = Q(L_R^i) - Q(L_R^j)$ and $Q_{ij}^L = Q(L_L^j) - Q(L_L^i)$.

Within the LRSM the Dirac mass matrices are hermitian. Therefore,

$$|Q(L_R^i) - Q(L_L^j) + Q(\phi_1)| = |Q(L_R^j) - Q(L_L^i) + Q(\phi_1)|, \quad (30)$$

with $i = 1, 2, 3$. In the appendix we show that since in our framework the Dirac mass matrices are both hermitian and hierarchical then the following relation is obtained:

$$Q_{ij}^R = Q_{ij}^L \equiv Q_{ij}. \quad (31)$$

In addition, from eq. (27) and eq. (28) we learn that in order to obtain neutrino masses below the eV scale, we must have:

$$m_{\nu_i}^D \lesssim m_\tau/100, \quad (32)$$

with $m_{\nu_i}^D$, $i = 1, 2, 3$ are the eigenvalues of the neutrino Dirac mass matrix.

Within our model, the suppression of the neutrino Dirac masses is achieved by aligning the sign of the U(1) charges of the lepton doublets with the one of ϕ_1 . This yields a suppression of $\varepsilon^{2|Q(\phi_1)|}$ of the neutrinos Dirac masses compared with the charged lepton ones. This is demonstrated in eqs. (24,25,27), that uses, $\text{sign}[Q(L_R^3) - Q(L_L^3)] = \text{sign}[Q(\phi_1)]$. Though a charge assignment in which $\text{sign}(Q_3) \neq \text{sign}(Q_{1,2})$ can, in principle, lead to a viable (hermitian and hierarchical) charged lepton mass matrix, it yields an enhancement of $\varepsilon^{-2|Q(\phi_1)|} = \mathcal{O}(100)$ to the corresponding entries in the neutrino mass matrix. It is enough that the first generation is assigned an opposite charge, $\text{sign}(Q_3) = \text{sign}(Q_2) \neq \text{sign}(Q_1)$, to typically have $m_{\nu_1}^D$ of the order of 100 MeV. This leads to a violation of the constraint in eq. (32). The same clearly holds for the other generations. Thus we must have:

$$\text{sign}(Q_1) = \text{sign}(Q_2) = \text{sign}(Q_3). \quad (33)$$

The hierarchy in the charged lepton masses and eqs. (30,31,33) lead to the following relations [17]:

$$\begin{aligned} \left| \frac{(M_\nu^D)_{12}}{(M_\nu^D)_{22}} \right| &\sim \left| \frac{M_{12}^{cl}}{M_{22}^{cl}} \right| \sim \sqrt{\frac{m_e}{m_\mu}}, & \left| \frac{(M_\nu^D)_{23}}{(M_\nu^D)_{33}} \right| &\sim \left| \frac{M_{23}^{cl}}{M_{33}^{cl}} \right| \sim \sqrt{\frac{m_\mu}{m_\tau}}, \\ \left| \frac{(M_\nu^D)_{13}}{(M_\nu^D)_{33}} \right| &\sim \left| \frac{M_{13}^{cl}}{M_{33}^{cl}} \right| \sim \sqrt{\frac{m_e}{m_\tau}}. \end{aligned} \quad (34)$$

From eq. (34) we learn that the charged lepton mass matrix typically gives very small mixing angles in the (12) and (13) planes. Thus LMA for the SN can only come from the light neutrino mass matrix.

With horizontal U(1) symmetries there are two typical structures of neutrino mass matrix that may account for both LMA in the (12) and (23) planes [18,19]. One of them involves a special structure of the 3×3 mass matrix. This structure cannot be reduced to a 2×2 block matrix description. In the appendix we show that in our case this type cannot lead to LMA in the (12) plane. In the second structure, the LMA of the SN comes from the entries of the 2×2 block mass matrix related to the first two generations. Below we separately investigate $M_{Se\bar{e}}^\nu$ and M_L^{Maj} and show that LMA in the relevant 2×2 block of the mass matrix cannot be obtained in our framework.

A. Mixing in M_{See}^ν

Consider the relevant 2×2 block in M_{See}^ν . Requiring a LMA between the first two generations is translated to the following condition:

$$(M_{See}^\nu)_{12} \gtrsim \max[(M_{See}^\nu)_{11}, (M_{See}^\nu)_{22}] . \quad (35)$$

Since the neutrino Dirac mass matrix, M_ν^D , is quasi diagonal it is clear that, to satisfy the condition of eq. (35), the following ratio between the entries of $(M_R^{Maj})^{-1}$ should hold:

$$R_{12}^R \equiv \left| \frac{(M_R^{Maj})_{22}^{-1}}{(M_R^{Maj})_{12}^{-1}} \right| \lesssim \frac{m_e}{m_\mu} . \quad (36)$$

In the appendix we show that the ratio R_{12}^R is larger than $\sqrt{\frac{m_e}{m_\mu}}$ and the condition of eq. (36) cannot be satisfied. Therefore LMA in the relevant block of M_{See}^ν is impossible. The above conclusion can be explained as follows. Within our framework strong hierarchy in the neutrino Dirac matrices, required by the hierarchy of the charged lepton masses, is inevitable. Such strong hierarchy suppresses any off-diagonal dominance that can be obtained in $(M_R^{Maj})^{-1}$. Therefore order one mixing in M_{See}^ν cannot be obtained.

B. Mixing in M_L^{Maj}

M_L^{Maj} is given by:

$$M_L^{Maj} = v_L \begin{pmatrix} \varepsilon^{2|Q'_3+Q_{13}|} & \varepsilon^{2|Q'_3+Q_{23}+Q_{13}|} & \varepsilon^{2|Q'_3+Q_{13}|} \\ \varepsilon^{2|2Q'_3+Q_{23}+Q_{13}|} & \varepsilon^{2|Q'_3+Q_{23}|} & \varepsilon^{2|2Q'_3+Q_{23}|} \\ \varepsilon^{2|2Q'_3+Q_{13}|} & \varepsilon^{2|2Q'_3+Q_{23}|} & \varepsilon^{2|Q'_3|} \end{pmatrix} , \quad (37)$$

where $2Q'_3 = 2Q(L_R^3) + Q(\Delta_R)$ and we used the fact that within LRSM $|M_L^{Maj}/v_L| = |M_R^{Maj}/v_R|$.

It is evident from eq. (37) that a proper charge assignment for $Q(\Delta_{R,L})$,

$$Q(\Delta_R) \simeq -2Q(L_R^3) - Q_{23} - Q_{13} ; \quad Q(\Delta_L) \simeq -2Q(L_L^3) + Q_{23} + Q_{13} , \quad (38)$$

yields dominance of $(M_L^{Maj})_{12}$ over the diagonal entries $(M_L^{Maj})_{11}$ and $(M_L^{Maj})_{22}$. However, the condition for large mixing, $(M_L^\nu)_{12} \gtrsim \max[(M_L^\nu)_{11}, (M_L^\nu)_{22}]$, together with eq. (7) imply that one needs also to compare $(M_L^\nu)_{12}$ to $(M_{See}^\nu)_{11}$ and $(M_{See}^\nu)_{22}$. As shown in eq. (22), the scales of M_L^{Maj} and M_{See}^ν are not independent in the LRSM. In particular, the higher the charges chosen for the triplet fields the smaller the product $v_L v_R$ becomes. Thus for a fixed value of v_R the contribution of M_L^{Maj} to M_l^ν might become negligible.

In our framework we saw that $(M_{See}^\nu)_{22}$ is expected to be dominant compared with the other entries of $(M_{See}^\nu)_{ij}$, $i, j = 1, 2$. Therefore we compare it to $(M_L^{Maj})_{12}$ assuming that the condition of eq. (38) holds. Given eq. (38) the determinant of M_R^{Maj} is given by:

$$\det(M_R^{Maj}) \sim v_R^3 \varepsilon^{|Q_{13}+Q_{23}|} . \quad (39)$$

Using eqs. (7,A4,39) we find that:

$$(M_{See}^\nu)_{22} \sim \frac{(m_{\nu_\mu}^D)^2}{v_R \varepsilon^{|Q_{12}|}} . \quad (40)$$

Hence, the ratio $(M_L^{Maj})_{12}/(M_{See}^\nu)_{22}$ is given by:

$$R_{12}^L \equiv (M_L^{Maj})_{12}/(M_{See}^\nu)_{22} \approx \frac{v_R v_L}{(m_{\nu_\tau}^D)^2} \varepsilon^{|Q_{12}|-4|Q_{23}|} . \quad (41)$$

Thus, LMA is induced by M_L^{Maj} if,

$$R_{12}^L \gtrsim 1 . \quad (42)$$

In the appendix we show that, for the ratio R_{12}^L to be larger than unity, it is required that $v_R \gtrsim 10^9$ which is irrelevant for our discussion. Therefore obtaining LMA in the (12) block is impossible.

In other words, we saw that in order to produce LMA in M_L^{Maj} one is driven to assign relatively high charges for $\Delta_{L,R}$ as shown in eq. (38). The high charges of $\Delta_{L,R}$ induce suppression of the product $v_L v_R$, as seen in eq. (22) and eq. (23). This suppression in its turn yields suppression of the ratio R_{12}^L .

C. Conclusion

From the above analysis we learn that a MLRSM with the additional horizontal U(1) symmetry discussed above cannot account for the LMA solution of the SN problem in a natural way. We focus on this point since the combined data from the SNO and Super-Kamiokande experiments [6,7] disfavors the SMA solution of the SN problem [8]. Though we do not show it explicitly in the text, a proper charge assignment for the various fields, *e.g.*, $Q(\Delta_L) = -Q(\Delta_R) = 8$, $Q(\phi_1) = 2$, $Q(L_L^{1,2,3}) = -Q(L_R^{1,2,3}) = 3, 4, 6$, and an appropriate choice of the ratio ε^4/r can produce a viable model of the lepton sector without any fine tuning. Such a model yields order one mixing for the atmospheric neutrinos and the SMA solution to the SN problem.

V. DISCUSSION AND CONCLUSION

A generic LRSM with the left-right symmetry (LRS) breaking scale $v_R \lesssim 15$ TeV and natural value for Yukawa Higgs couplings leads to a severe problem in the lepton sector: The light neutrino masses are in fact rather heavy. The problem arises from two different,

generically, independent parts of the model:

1. v_L , the triplet VEV is relatively high, $v_L \sim k^2/v_R = \mathcal{O}(1 \text{ GeV})$, which leads to a large eigen values of the Majorana mass matrix for the left-handed neutrinos.
2. The scale of the neutrino mass matrix obtained by the seesaw mechanism is large, with the mass of the heaviest neutrino of the order $m_\tau^2/v_R = 0.1 \text{ MeV}$, provided that the tau neutrino Yukawa couplings are similar to those of the tau.

In this context we demonstrated that, within a minimal LRSM, the two problems are solved using an additional U(1) symmetry, when the ratio between the two VEVs of the bi-doublet neutral fields is assumed to be small, of the order $10^{-2} - 10^{-3}$.

We further showed that, if we assume that the same U(1) symmetry explains the charged lepton flavor hierarchy, small mixing angle is obtained between the first two generations. Nevertheless, there is no reason that a viable model, which combines the above U(1) symmetry with additional symmetries, cannot be constructed.

This work is motivated by two reasons: First, the idea that the fundamental scale of gravity might be much smaller than the Planck scale was recently proposed and is, at present, consistent with all of our experimental knowledge [20]. Our work demonstrates how a LRSM, in which the fundamental scale of gravity (and other NP sources) is relatively low, can account in a natural way for the smallness of neutrino masses. To the best of our knowledge no such complete LRSM was considered so far in the literature [21].

Our second motivation is related to the fact that LRSM with spontaneous CP violation (SCPV) [9,13] were recently analyzed in the literature [10,11,16,22–26] and found to have a considerable agreement with all the CP conserving experimental data. It was also shown there that constraints from the quark sector typically require r to be of $\mathcal{O}(10^{-2} - 10^{-3})$ [11,23,25,26], and that v_R should be of $\mathcal{O}(10 \text{ TeV})$ [11,25].

In that context it is important to note that MLRSM with SCPV cannot account for the observed CPV in the quark sector [13,15,16]. Recently it was also shown in [16] that the CP properties of the vacuum of the MLRSM with SCPV are strongly connected to the physical Higgs fields mass spectrum. This means that the above problem of the MLRSM with SCPV might be shared by many LRSM with SCPV. To what extent this problem is really general is yet to be investigated. Therefore LRSM with SCPV should not be discarded.

Finally we remark that the recent result of BaBar [27] and Belle [28] experiments exclude any LRSM with SCPV in which the quarks fields couples only to a single bidoublet field [11,25,26,29].

ACKNOWLEDGMENTS

We thank Yossi Nir for helpful discussions and comments on the manuscript.

APPENDIX A:

1. The Values of Q_{ij}^{LR}

In this subsection we prove eq. (31), that is, $Q_{ij}^R = Q_{ij}^L$. Consider a two generation picture and suppose that $Q_{12}^R \neq Q_{12}^L$ and that, without loss of generality, $\text{sign}(Q_{12}^R) = \text{sign}(Q_2)$, with $Q_2 = Q(L_R^2) - Q(L_L^2) + Q(\phi_1) > 0$. The only way to ensure that the product $M_{12} \cdot M_{21}$ is of the order of $m_1 m_2$, as required by the hierarchy of the charged lepton masses, is when Q_{12}^L is given by:

$$Q_{12}^L \simeq -(Q_{12}^R + 2Q_2). \quad (\text{A1})$$

In this case it is easy to see that M_{11} is of the same order as M_{22} which does not allow for $m_1 \ll m_2$. Therefore we conclude that eq. (31) must be satisfied.

2. Mixing in a Non-reducible 3×3 Majorana Neutrino Mass Matrix

In ref [18] it was shown how an $L_e - L_\mu - L_\tau$ symmetry can lead to LMA both for the AN and for the SN. However it was assumed in [18] that the hierarchy in the charged lepton masses comes from additional flavor symmetry. Furthermore the 2×2 block related to the second and third generations in the Dirac neutrino mass matrix was assumed to be non-hierarchical. In our case we do not allow for additional continuous symmetries, therefore an $L_e - L_\mu - L_\tau$ symmetry cannot be realized, and the Dirac matrices obey the structure given in eq. (34). Nevertheless we want to consider the case that neutrino Majorana mass matrices have an approximate $L_e - L_\mu - L_\tau$ structure²:

$$M_{L,R}^{Maj} \approx v_{L,R} \begin{pmatrix} \varepsilon' & a & 1 \\ a & \varepsilon' & \varepsilon' \\ 1 & \varepsilon' & \varepsilon' \end{pmatrix}, \quad (\text{A2})$$

with $\varepsilon' \ll |a| \lesssim 1$.

Applying the analysis of [18] to our model and assuming for simplicity a diagonal Dirac mass matrix, typically yields the following structure for M_{See}^ν :

$$M_{See}^\nu \sim \frac{\varepsilon^{4|Q(\phi_1)|}}{v_R} \begin{pmatrix} \varepsilon' m_e^2 & m_\mu m_e & m_e m_\tau \\ m_\mu m_e & \varepsilon' m_\mu^2 & 0 \\ m_\tau m_e & 0 & \frac{m_\tau^2}{\varepsilon'} \end{pmatrix}, \quad (\text{A3})$$

²This can be achieved, *e.g.*, using the following charge assignment, $Q(\Delta_L) = -Q(\Delta_R) = 8.5$, $Q(\phi_1) = 2$, $Q(L_L^{1,2,3}) = -Q(L_R^{1,2,3}) = 3, 4, 11/2$, this leads to electron parity symmetry which we assume to be broken by another small parameter $\varepsilon' \ll \varepsilon$.

where we assumed also that $r \sim \varepsilon^{2|Q(\phi_1)|}$. As clearly seen from eq. (A3) to have LMA for the SN one needs $\varepsilon' \lesssim \frac{m_e}{m_\mu}$. This means that the heavy neutrino mass is enhanced by the inverse of the same factor which clearly violates eq. (32). In other words, LMA cannot be induced by M_{See}^ν .

In the case where ε' is relatively large, as required by eq. (32), LMA also cannot be induced by M_L^{Maj} . To see that one needs to verify that $\frac{(M_L^{Maj})_{12}}{(M_{See}^\nu)_{22}} = R_{12}^L \ll 1$. This ratio is investigated in subsection IV B and in subsection A 4, below, in a similar situation. Applying the same analysis to the present case one finds that the value of the ratio R_{12}^L is enhanced by a factor of $1/\varepsilon'$ relative to the corresponding values given in eqs. (A12,A13,A14). The mild enhancement, of $\mathcal{O}(1/\varepsilon')$, still yields $R_{12}^L \ll 1$ from which we learn that the mixing angle induced by M_L^{Maj} are small.

Thus the above scenario cannot lead to LMA between the first two generations.

3. Mixing in M_{See}^ν

Below we show that the ratio R_{12}^R defined in eq. (36) is always larger than $\sqrt{\frac{m_e}{m_\mu}}$ and therefore eq. (36) cannot be satisfied. In order to do so we focus on the structure of $(M_R^{Maj})^{-1}$,

$$(M_R^{Maj})^{-1} = \frac{1}{v_R} \begin{pmatrix} \varepsilon^{2|Q'_3+Q_{13}|} & \varepsilon^{|2Q'_3+Q_{23}+Q_{13}|} & \varepsilon^{|2Q'_3+Q_{13}|} \\ \varepsilon^{|2Q'_3+Q_{23}+Q_{13}|} & \varepsilon^{2|Q'_3+Q_{23}|} & \varepsilon^{|2Q'_3+Q_{23}|} \\ \varepsilon^{|2Q'_3+Q_{13}|} & \varepsilon^{|2Q'_3+Q_{23}|} & \varepsilon^{2|Q'_3|} \end{pmatrix}^{-1}. \quad (A4)$$

From eq. (A4) we find that the ratio R_{12}^R is given by:

$$R_{12}^R = \left| \frac{\varepsilon^{2|Q'_3+Q_{13}|+2|Q'_3|} - \varepsilon^{2|2Q'_3+Q_{13}|}}{\varepsilon^{|2Q'_3+Q_{13}+Q_{23}|+2|Q'_3|} - \varepsilon^{|2Q'_3+Q_{13}|+|2Q'_3+Q_{23}|}} \right|. \quad (A5)$$

To see whether eq. (36) can be satisfied we are interested in the minimal value that the ratio R_{12}^R can have. To find it we divide it into separate ranges of Q' and calculate the minimal value of R_{12}^R in each region. Without loss of generality we assume that Q' is non-negative, the result we get at the end of the discussion is valid for any value of Q' .

- $Q_{13} > -Q'$:

In that case all the terms in the exponents of eq. (A5) are positive and it is easy to see that R_{12}^R is given by:

$$R_{12}^R \sim \varepsilon^{|Q_{12}|} \sim \sqrt{\frac{m_e}{m_\mu}}, \quad (A6)$$

thus the condition of eq. (36) is not satisfied.

- $-2Q' \geq Q_{13} \geq -Q'$ and $-2Q' \geq Q_{13} + Q_{23}$:

In that case only the exponent of the first term in the dominator is negative and the ratio is given by:

$$R_{12}^R \sim \max \frac{[\varepsilon^{|2Q_{13}|}, \varepsilon^{4Q'_3+2Q_{13}}]}{\varepsilon^{4Q'_3+Q_{13}+Q_{23}}} \geq \varepsilon^{|Q_{12}|} \sim \sqrt{\frac{m_e}{m_\mu}}, \quad (\text{A7})$$

where in the last line we used the relation $Q_{13} = Q_{12} + Q_{23}$. Therefore, the condition of eq. (36) is not satisfied.

- $-2Q' \geq Q_{13} \geq -Q'$ and $-2Q' > Q_{13} + Q_{23}$:

In that case the exponent of the first term in the dominator and also that of the first term in the denominator are negative and the ratio is given by:

$$R_{12}^R \sim \frac{\max [\varepsilon^{|2Q_{13}|}, \varepsilon^{4Q'_3+2Q_{13}}]}{\max [\varepsilon^{2|Q_{13}+Q_{12}|}, \varepsilon^{4Q'_3+Q_{13}+Q_{23}}]} \geq \varepsilon^{|Q_{12}|} \sim \sqrt{\frac{m_e}{m_\mu}}, \quad (\text{A8})$$

and the condition of eq. (36) is not satisfied.

- $-2Q' > Q_{13}$ and $-2Q' \leq Q_{23}$:

In that case the exponent of the first term in the dominator and also that of the first term in the denominator are negative. Therefore R_{12}^R is given by:

$$R_{12}^R \sim \frac{\varepsilon^{|2Q_{13}|}}{\varepsilon^{2|Q_{13}+Q_{12}|}} \geq \varepsilon^{|Q_{12}|} \sim \sqrt{\frac{m_e}{m_\mu}}, \quad (\text{A9})$$

and the condition of eq. (36) is not satisfied.

- $-2Q' > Q_{13}$ and $-2Q' > Q_{23}$:

In that case the exponent of the first term in the dominator and also that of the first term in the denominator are negative. Therefore the ratio is given by:

$$R_{12}^R \sim \frac{\max [\varepsilon^{|2Q_{13}|}, \varepsilon^{-4Q'_3-2Q_{13}}]}{\max [\varepsilon^{2|Q_{13}+Q_{12}|}, \varepsilon^{-4Q'_3-Q_{13}-Q_{23}}]} \geq \varepsilon^{|Q_{12}|} \sim \sqrt{\frac{m_e}{m_\mu}}, \quad (\text{A10})$$

and the condition of eq. (36) is not satisfied.

Thus we conclude that mixing of order one in $M_{S_{ee}}''$ between the first two generation cannot be achieved.

4. Mixing in M_L^{Maj}

Below we show that the ratio R_{12}^L , defined in eq. (41), is larger than unity only if $v_R \gtrsim 10^9$ which is irrelevant for our discussion and therefore eq. (42) cannot be satisfied. The ratio R_{12}^L is given by:

$$\begin{aligned}
R_{12}^L &\approx \frac{v_R v_L}{(m_{\nu_\tau}^D)^2} \varepsilon^{|Q_{12}| - 4|Q_{23}|} \sim \frac{k_1^2}{(m_{\nu_\tau}^D)^2} \varepsilon^{|Q_{12}| - 4|Q_{23}|} \times \\
&\quad \max[r^2 \varepsilon^{2|Q_{23} + Q_{13} + Q_3 - Q(\phi_1)|}, r \varepsilon^{2|Q_{23} + Q_{13} + Q_3|}, \varepsilon^{2|Q_{23} + Q_{13} + Q_3 + Q(\phi_1)|}] \\
&\sim \varepsilon^{|Q_{12}| - 4|Q_{23}|} \frac{\max[r^2 \varepsilon^{2|Q_{23} + Q_{13} + Q_3 - Q(\phi_1)|}, r \varepsilon^{2|Q_{23} + Q_{13} + Q_3|}, \varepsilon^{2|Q_{23} + Q_{13} + Q_3 + Q(\phi_1)|}]}{\max[r^2 \varepsilon^{2|Q_3 + Q(\phi_1)|}, r \varepsilon^{|Q_3 - Q(\phi_1)| + |Q_3 + Q(\phi_1)|}, \varepsilon^{2|Q_3 - Q(\phi_1)|}]}, \quad (A11)
\end{aligned}$$

where in the second line we used eqs. (22,23,38) to simplify the above expression. To see whether the ratio R_{12}^L could be equal or larger than unity we investigate eq. (A11) in four regimes relates to $Q(\phi_1)$. We assume that Q_3 and also Q_{ij} are non-negative (since they carry the same sign according to our above assumption). This assumption does not affect our final conclusion since we do not impose any further assumptions on $Q(\phi_1)$:

(i) $Q_3 \geq Q(\phi_1) \geq 0$:

The ratio of eq. (A11) is given by,

$$R_{12}^L \sim \varepsilon^{3|Q_{12}|} \cdot \max[r^2, \varepsilon^{2Q(\phi_1)}, \varepsilon^{4Q(\phi_1)}] \ll 1, \quad (A12)$$

which means that in this case no large mixing is possible.

(ii) $Q_{23} + Q_{13} + Q_3 \geq Q(\phi_1) > Q_3$:

The ratio of eq. (A11) is given by,

$$R_{12}^L \sim \varepsilon^{3|Q_{12}|} \varepsilon^{4Q_3} \ll 1, \quad (A13)$$

which means that in this case no large mixing is possible.

(iii) $Q(\phi_1) > Q_{23} + Q_{13} + Q_3$:

The ratio of eq. (A11) is given by:

$$R_{12}^L \lesssim \varepsilon^{3|Q_{12}|} \varepsilon^{|Q_{23} + Q_{13}|} \ll 1, \quad (A14)$$

where we used the fact that we require $r \lesssim \varepsilon^{2|Q(\phi_1)|}$ as discussed in section III. Eq. (A14) implies that no large mixing is possible.

- (iv) In items (i)-(iii) we considered cases with $Q(\phi_1) > 0$. Let us consider the case with negative charges for example consider the range $-Q_3 < Q(\phi_1) < 0$:

The ratio of eq. (A11) is given by,

$$R_{12}^L \sim \varepsilon^{3|Q_{12}|} r^{-2}, \quad (\text{A15})$$

From eq. (A15) we learn that maximal mixing is possible if

$$r \lesssim \varepsilon^{1.5|Q_{12}|} \sim 1/50, \quad (\text{A16})$$

which means that large mixing is in principle possible.

However, as discussed above, opposite charges of ϕ and Q_3 yield inverse hierarchy between the neutrino and charge lepton Dirac masses,

$$\frac{m_{\nu_i}^D}{m_i^D} \sim \varepsilon^{-2|Q\phi|} \gtrsim 1. \quad (\text{A17})$$

It means that in order not to produce too large neutrino masses v_R is bounded from below:

$$v_R \geq \frac{(m_D^{\nu\tau})^2}{1 \text{ eV}} \gtrsim \frac{m_\tau^2}{1 \text{ eV}} \sim 10^9 \text{ GeV}. \quad (\text{A18})$$

Thus in this case large mixing cannot be obtained with a low value of v_R . A similar conclusion is clearly obtained for any negative value of $Q(\phi_1)$.

REFERENCES

- [1] J.C. Pati and A. Salam, Phys. Rev. **D 10** (1974) 275; R.N. Mohapatra and J.C. Pati, Phys. Rev. **D 11** (1975) 566 and 2558; G. Senjanović and R.N. Mohapatra, Phys. Rev. **D 12** (1975) 1502; G. Senjanović, Nucl. Phys. **B153** (1979) 334; R.N. Mohapatra and G. Senjanović, Phys. Rev. Lett. **44** (1980) 912; Phys. Rev. **D 23** (1981) 165; C. S. Lim and T. Inami, Prog. Theor. Phys. **67** (1982) 1569.
- [2] T. G. Rizzo and G. Senjanović, Phys. Rev. Lett. **46** (1981) 1315; Phys. Rev. **D24** (1981) 704; Phys. Rev. **D 25** (1982) 235; M. Fukugita, T. Yanagida and M. Yoshimura, Phys. Lett. **B106** (1981) 183; R. N. Mohapatra, Fortsch. Phys. **31** (1983) 185.
- [3] S.S. Gershtein and Ya. B. Zeldovich, JETP Lett. **4** (1966) 120; R. Cowsik and J. McClelland, Phys. Rev. Lett. **29** (1972) 669; J.R. Primack and M.A.K. Gross, [astro-ph/0007165](#), to appear in *Current Aspects of Neutrino Physics*, ed. D. O. Caldwell (Springer, Berlin Heidelberg 2000).
- [4] D.E. Groom *et al.*, Eur. Phys. J. **C 15** (2000) 1.
- [5] L. Baudis *et al.*, Heidelberg-Moscow Double-Beta Decay Exp., Phys. Rev. Lett. **83** (1999) 41 [[hep-ex/9902014](#)]; H.V. Klapdor-Kleingrothaus *et al.*, Heidelberg-Moscow Double-Beta Decay Exp., Talk presented at the third International Conference- DARK2000, to be published. in Proc. of DARK2000, Springer (2000), [hep-ph/0103062](#).
- [6] Q.R. Ahmad *et al.*, SNO Collaboration, Phys. Rev. Lett. **87**, 071301 (2001) [[astro-ph/0106015](#)].
- [7] B.T. Cleveland *et al.*, Astrophys. J. **496** (1998) 505; W. Hampel *et al.*, GALLEX Collaboration, Phys. Lett. **B447** (1999) 127; J.N. Abdurashitov *et al.*, SAGE Collaboration, Phys. Rev. **C 60** (1999) 055801 [[astro-ph/9907113](#)]; M. Altmann *et al.*, GNO Collaboration, Phys. Lett. **B490** (2000) 16 [[hep-ex/0006034](#)]; S. Fukuda *et al.*, Super-Kamiokande Collaboration, Phys. Rev. Lett. **81** (1988) 4279; Phys. Rev. Lett. **86** (2001) 5651; M. Apollonio *et al.*, CHOOZ Collaboration, Phys. Lett. **B466** (1999) 415 [[hep-ex/9907037](#)].
- [8] A. Bandyopadhyay, S. Choubey, S. Goswami and Kamales Kar [hep-ph/0106264](#); J. N. Bahcall, M.C. Gonzalez-Garcia and C. Pena-Garay, [hep-ph/0106258](#); V. Barger, D. Marfatia and K. Whisnant [hep-ph/0106207](#); G.L. Fogli, E. Lisi, D. Montanino and A. Palazzo, [hep-ph/0106247](#).
- [9] G. Ecker, W. Grimus and H. Neufeld, Nucl. Phys. **B 247** (1984) 70; G. Ecker and W. Grimus, Nucl. Phys. **B 258** (1985) 328; Z. Phys. **C30** (1986) 293; D. Chang, Nucl.

- Phys. **B 214** (1983) 435; J. M. Frere *et. al.*, Phys. Rev. **D 46** (1992) 337; J. Basecq and D. Wyler, Phys. Rev. **D39** (1989) 870; G.C. Branco and L. Lavoura, Phys. Lett. **B165** (1985) 327.
- [10] G. Barenboim, J. Bernabeu and M. Raidal, Nucl. Phys. **B478** (1996) 527 [hep-ph/9608450]; Z. Phys. C **73** (1997) 321 [hep-ph/9603379].
- [11] P. Ball, J.M. Frere and J. Matias, Nucl. Phys. **B 572** (2000) 3 [hep-ph/9910211].
- [12] M. Gell-Mann, P. Ramond and R. Slansky, in *Supergravity*, ed. P. van Nieuwenhuizen and D. Freedman (North-Holland 1979); T. Yanagida, in *Proceedings of the Workshop on the Unified Theory and the Baryon Number in the Universe*, ed. O. Sawada and A. Sugamoto (Tsukuba 1979); R. N. Mohapatra and G. Senjanović, Phys. Rev. Lett. **44**, 912 (1980).
- [13] J. Basecq, J. Liu, J. Milutinovic and L. Wolfenstein, Nucl. Phys. **B272** (1986) 145.
- [14] J.F. Gunion, J. Grifols, A. Mendez, B. Kayser and F. Olness, Phys. Rev. **D 40** (1989) 1546.
- [15] N.G. Deshpande, J.F. Gunion, B. Kayser and F. Olness, Phys. Rev. **D 44**, 837 (1991).
- [16] G. Barenboim, M. Gorbahn, U. Nierste and M. Raidal, hep-ph/0107121.
- [17] Y Grossman and Y. Nir, Nucl. Phys. **B448** (1995) 30 [hep-ph/9502418].
- [18] R. Barbieri *et. al.*, JHEP **12** (1998) 98017 [hep-ph/9807235].
- [19] Y. Grossman, Y Nir and Y. Shadmi, JHEP **9810** (1998) 007 [hep-ph/9808355].
- [20] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. **B429** (1998) 263 [hep-ph/9803315]; I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. **B436** (1998) 257 [hep-ph/9804398]; N. Arkani-Hamed and S. Dimopoulos hep-ph/9811353; Z. Berezhiani and Gia Dvali, Phys. Lett. **B450** (1999) 24 [hep-ph/9811378]; N. Arkani-Hamed, L. Hall, D. Smith and N. Weiner, Phys. Rev. **D61** (2000) 116003 [hep-ph/9909326].
- [21] Papers that discuss small neutrino masses induced by a low-energy seesaw like mechanism in a non-LRS framework: E. Ma, Phys. Rev. Lett. **86** (2001) 2502 [hep-ph/0011121]; hep-ph/0107177; hep-ph/0103278; E. Ma, M. Raidal and U. Sarkar, hep-ph/0012101; E. Ma and M. Raidal, hep-ph/0012366; M.B. Tully and G.C. Joshi, Phys. Rev. **D64** (2001) 011301 [hep-ph/0011172]; Z. Tavartkiladze, hep-ph/0105281.
- [22] G. Barenboim *et. al.*, Nucl. Phys. **B511** (1998) 577 [hep-ph/9611347].

- [23] G. Barenboim, J. Bernabeu and M. Raidal, Phys. Rev. **D 55** (1997) 4213 [hep-ph/9702337].
- [24] G. Barenboim, J. Bernabeu, J. Matias and M. Raidal, Phys. Rev. **D60** (1999) 016003 [hep-ph/9901265].
- [25] P. Ball and R. Fleischer, Phys. Lett. **B 475** (2000) 111 [hep-ph/9912319].
- [26] S. Bergmann and G. Perez, hep-ph/0103299, Phys. Rev. **D**, in press.
- [27] B. Aubert *et al.*, BaBar collaboration, hep-ex/0107013.
- [28] K. Abe *et al.*, BELLE collaboration, hep-ex/0107061.
- [29] M. Beneke, Plenary Talk, the EPS Int. Conference on High Energy Physics, July 14 (2001-Budapest).